

C & C Peat Co., Inc.

Premium Quality Potting Media

(800) 330-4866



<http://www.ccpeat.com>

info@ccpeat.com

Technical Information for C&C Peat Company's compost

As a key component in Enviro-Peat and other potting media mixes

C&C Peat is currently composting materials taken primarily from many horse farms in Marion County, as well as secondarily from land clearing debris. This material is brought into our fully licensed composting facility where it will undergo a rigorous series of treatments to be transformed into a useful product for plant growth. To ensure quality control we are restricting our feedstock source. It is difficult to determine what pesticides or other chemicals might be in home yard debris, so we are not taking this material as part of our compost feedstock. Because of this control, we have the ability to maintain a consistent level of nutrient content.

After entering our facility, the material to be composted is placed in long windrows. These windrows will be treated with moisture, as is needed, in order to maintain aerobic composting. Through the first 15 days, each windrow will be turned at least 5 times; the compost is then stabilized for 75 days and cured for an additional 30 days. This allows the windrows to maintain higher oxygen content, reducing odors and assisting beneficial microbes. Through the turning process, the compost will maintain a temperature of at least 131 degrees Fahrenheit, reaching a peak temperature of 155 degrees Fahrenheit before turning. This is accomplished through the decomposition of the feeder material.

Pathogenic organisms such as bacteria, viruses, fungi, and protozoa are found in small or background concentrations with other living organisms everywhere on earth. The concern for human, animal, and plant health from pathogens comes when these organisms are found in large concentrations. Larger concentrations of pathogens are found in animal and pet manures and biosolids. When these materials are used as a feedstock, the following standards must be followed to prevent a health risk.

The EPA in Part 503 sets standards for the destruction and concentration of pathogens in composted biosolids. The standards set in Part 503 are often used as marketplace specifications for compost or soil amendment sold for public use. By maintaining these standards, the following harmful pathogens are destroyed:

Pathogen	Temperature at Time of Pathogen Death
Salmonella typhosa	No growth beyond 115°F, death within 30 minutes at 131-140°F, or within 20 minutes beyond 140°F
Salmonella sp.	Death within 1 hour at 131°F or within 15-20 minutes at 140°F
Shigella sp	Death within 1 hour at 131°F.
Escherichia coli (E. coli)	Most die within 1 hour at 130°F or all within 15-20 minutes at 140°F.
Entamoeba histolytica cysts	Death within a few minutes at 113°F or within a few seconds at 131°F.
Taenia Saginata	Death within a few minutes at 131°F.
Trichinella spiralis larvae	Death at 131°F.
Brucella abortus	Death within 3 minutes at 147-145°F or within 1 hour at 131°F.
Micrococcus pyogenes var. hominis	Death within 10 minutes at 122°F.
Streptococcus pyogenes	Death within 10 minutes at 130°F.
Mycobacterium tuberculosis var. hominis	Death within 15-20 minutes at 151°F or momentarily at 153°F.
Corynebacterium diphtheria	Death within 45 minutes at 131°F.
Necator americanus	Death within 50 minutes at 113°F.
Ascaris lumbricoides eggs	Death in less than 1 hour at temperatures over 122°F

In the first 15 days of composting (the "thermophilic" phase) temperatures reach as high as 155 degrees, but never lower than 131 degrees, resulting in all weed seeds being destroyed. Many of these seeds were present as the result of being accidentally incorporated in the feed given to horses, as well as from windblown sources. Once the minimum temperature of 131 degrees is reached, these seeds are no longer a problem. (<http://www.cias.wisc.edu/pdf/artofcompost.pdf>)

In addition to the reduction of potentially harmful pathogens and weed seeds, the composting process allows for the control of heavy metal concentrations that must remain below a regulated level, per IAC105.3 (455B, 455D). The specific heavy metal maximum concentrations permitted are:

	Arsenic	Cadmium	Copper	Lead	Mercury	Nickel	Selenium	Zinc
Parts per million	41	39	1500	300	17	420	36	2800

Fertilizers



Quality Potting Media



Quality Service

1650 County Road 470

Okahumpka, FL 34762

Office: (352) 323-8213

Fax: (352) 365-0367